



Calf Starter with Decco

For Dairy and Beef Calves on Milk.

MEDICATED

For the prevention of coccidiosis in ruminating and non-ruminating calves and cattle caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Decoquinat 17.8 mg/lb (0.0039 %)

Guaranteed Analysis

| | | |
|------------------------------|------|---|
| Crude Protein ,Not Less than | 17.0 | % |
| Crude Fat ,Not Less than | 2.8 | % |
| Crude Fiber ,Not More than | 6.8 | % |
| Calcium ,Not Less than | 0.4 | % |
| Calcium ,Not More than | 0.8 | % |
| Phosphorus ,Not Less than | 0.2 | % |
| Ash ,Not More than | 6 | % |

Ingredients

Rolled Corn, Soybean Meal, Rolled Oats, Rolled Barley, Wheat Mill Run, Cane Molasses, Ground Corn, Limestone, Rice Bran, Salt, Flavor, Vitamin E, Calcium Carbonate, Choline Chloride, Iron Sulfate, Zinc Oxide, Manganese Oxide, Niacin, D-Calcium Pantothenate, Copper Sulfate, Vitamin A, D, E supplement, Riboflavin Supplement, Sodium Selenite, Vitamin K supplement, Pyridoxine HCl, Vitamin B-12, Thiamin, Folic Acid, Ethylenediamine Dihydroiodide, Cobalt Carbonate, Sodium Bicarbonate, Zinc Sulfate, Manganous Sulfate, Magnesium Oxide, Calcium Chloride, Vitamin E supplement and Vitamin D3.

Feeding Directions

Feed to Calves from 3 days of age to weaning. Provide clean fresh water at all times.

Warning

WARNING: Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed type C feed or milk replacer to provide 22.7 mg per 100 lb. bodyweight (0.5 mg/kg) per day.

CAUTION: Do not use in feeds containing bentonite.

Manufactured By:
Hunt & Behrens, Inc.
30 Lakeville St.
Petaluma, CA 94952
(707) 762-4594

50 pounds

Feed Formula Number:

1610

MEDICATED

Calf Grower with Decco

For weaned dairy and beef calves up to 400 pounds.

MEDICATED

For the prevention of coccidiosis in ruminating and non-ruminating calves and cattle caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Decoquinat 14.1 mg/lb (0.0031 %)

Guaranteed Analysis

| | | |
|------------------------------|------|---|
| Crude Protein ,Not Less than | 15.0 | % |
| Crude Fat ,Not Less than | 2.8 | % |
| Crude Fiber ,Not More than | 6.2 | % |
| Calcium ,Not Less than | 0.2 | % |
| Calcium ,Not More than | 0.6 | % |
| Phosphorus ,Not Less than | 0.2 | % |
| Ash ,Not More than | 5 | % |

Ingredients

Rolled Corn, Soybean Meal, Rolled Barley, Rolled Oats, Wheat Mill Run, Cane Molasses, Ground Corn, Limestone, Rice Bran, Salt, Flavor, Vitamin E, Calcium Carbonate, Choline Chloride, Iron Sulfate, Zinc Oxide, Manganese Oxide, Niacin, D-Calcium Pantothenate, Copper Sulfate, Vitamin A, D, E supplement, Riboflavin Supplement, Sodium Selenite, Vitamin K supplement, Pyridoxine HCl, Vitamin B-12, Thiamin, Folic Acid, Ethylenediamine Dihydroiodide, Cobalt Carbonate, Sodium Bicarbonate, Zinc Sulfate, Manganous Sulfate, Magnesium Oxide, Calcium Chloride, Vitamin E supplement and Vitamin D3.

Feeding Directions

Feed to weaned calves consuming at least 2 pounds of grain per day. Provide clean fresh water at all times.

Warning

WARNING: Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed type C feed or milk replacer to provide 22.7 mg per 100 lb. bodyweight (0.5 mg/kg) per day.

CAUTION: Do not use in feeds containing bentonite.

Manufactured By:
Hunt & Behrens, Inc.
30 Lakeville St.
Petaluma, CA 94952
(707) 762-4594

50 pounds

Feed Formula Number:

1606

MEDICATED