



**Show Goat
With Grand Goat™ and Full Stream™**

For Growing Goats

MEDICATED

For the prevention of coccidiosis in young goats caused by Eimeria christenseni and Eimeria ninakohlyakimovae. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Decoquinatate 13.6 g/ton

Guaranteed Analysis

| | |
|--|--------|
| Crude Protein* ,Not Less than | 16.8 % |
| * Contains not more than 0.39 % crude protein from Non-protein Nitrogen. | |
| Crude Fat ,Not Less than | 3.8 % |
| Crude Fiber ,Not More than | 12.0 % |
| Calcium ,Not Less than | 0.8 % |
| Calcium ,Not More than | 1.2 % |
| Phosphorus ,Not Less than | 0.4 % |
| Ash ,Not More than | 9 % |

Ingredients

Wheat Mill Run, Ground Grains, Oat Hay Meal, Corn Distillers Grains, Soybean Meal, Alfalfa Meal, Cane Molasses, Rice Bran, Limestone, Salt, Ammonium Chloride, Flavor, Vit. A supplement, Sodium Bicarbonate, Zinc Sulfate, Manganous Sulfate, Magnesium Oxide, Calcium Chloride, Vitamin E supplement, Copper Sulfate, Sodium Selenite, Vitamin A, Iron Sulfate, Vitamin D3, Ethylenediamine Dihydroiodide, Cobalt Carbonate, Wheat Germ Meal, Brewers Dried Yeast, Zinc methionine complex, Dried Potato products, Calcium citrate, plant protein products, Vit D3 supplement and Vit E supplement.

Feeding Directions

Feed at a rate of 1-2% of Body weight daily.
Always provide clean fresh water, salt, and adequate forage

Warning

WARNING: Do not feed to goats producing milk for food.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard
Contains 30.87 ppm Copper. Do not feed to sheep
or other Copper sensitive species.

Manufactured By:



Hunt & Behrens, Inc.
30 Lakeville St.
Petaluma, CA 94952
(707) 762-4594

Feed Formula Number 754

MEDICATED

Net Weight 50 pounds

Goat Feeding Guidelines

- Always provide plenty of clean, fresh water.
- Make feed changes gradually. Increase feed amounts over 3-5 days.
- Feed at least twice a day.
- Weigh your feed. At least know what a full measure weighs.
- Treat your goat for internal and external parasites according to manufacturer directions. Always observe product withdrawal times.
- Feed is perishable. Keep in a clean dry place. Use within four weeks of purchase.
- Weigh your goat periodically to track rate of gain and time to target weight. Adjust feeding amounts to achieve optimum body condition.
- Practice showing your goat regularly. Not only does this make you a better showman, the exercise improves your goat's condition.



Hunt & Behrens, Inc.
30 Lakeville St.
Petaluma, CA 94952
(707) 762-4594